

Local Controllability of some PDEs

In collaboration with Jean-Michel Coron and Hoai-Minh Nguyen

Armand Koenig

18 January 2023

Conférence Itinérante du GDR AEDP

Introduction

Definition

A control system is an equation of the form

$$\dot{X}(t) = f(X(t), u(t))$$

- State space H
- Control space U
- $f: H \times U \rightarrow H$
- State $X: [0, T] \rightarrow H$
- Control $u: [0, T] \rightarrow U$

Definition

A control system is an equation of the form

$$\dot{X}(t) = f(X(t), u(t))$$

- State space H
- Control space U
- $f: H \times U \rightarrow H$
- State $X: [0, T] \rightarrow H$
- Control $u: [0, T] \rightarrow U$

Control theory

- Range of $u \mapsto X(T)$?
- For every $X_0, \exists u$ such that $X(T) = 0$?


Small-time Local Controllability (around 0)

$\dot{X} = f(X, u)$ with $f(0, 0) = 0$.

0 •


Small-time Local Controllability (around 0)

$\dot{X} = f(X, u)$ with $f(0, 0) = 0$. For $\epsilon > 0$




Small-time Local Controllability (around 0)

$\dot{X} = f(X, u)$ with $f(0, 0) = 0$. For $\epsilon > 0$, does there exist $\eta > 0$




Small-time Local Controllability (around 0)


$\dot{X} = f(X, u)$ with $f(0, 0) = 0$. For $\epsilon > 0$, does there exist $\eta > 0$ such that if $0 < T < \epsilon$, $|X_0| < \eta$, $|X_T| < \eta$



Small-time Local Controllability (around 0)
 $\dot{X} = f(X, u)$ with $f(0, 0) = 0$. For $\epsilon > 0$, does there exist $\eta > 0$ such that if $0 < T < \epsilon$, $|X_0| < \eta$, $|X_T| < \eta$, we can find $|u|_{L^\infty(0, T)} < \epsilon$ such that $X(T) = X_T$?



Small-time Local Controllability (around 0)
 $\dot{X} = f(X, u)$ with $f(0, 0) = 0$. For $\epsilon > 0$, does there exist $\eta > 0$ such that if $0 < T < \epsilon$, $|X_0| < \eta$, $|X_T| < \eta$, we can find $|u|_{L^\infty(0, T)} < \epsilon$ such that $X(T) = X_T$?



Theorem (Linear test)

Small-time local controllability holds if the linearized equation is controllable.

Proof.

$$\dot{X} = L_1 X + L_2 u + NL(X, u)$$

$$\mathcal{F}: g \mapsto Y \text{ solution to } \dot{Y} = L_1 Y + g, Y(0) = 0$$

Banach fixed-point theorem to $(X, u) \mapsto (Y, v)$ where

$$\begin{cases} v := \text{Linear control}(X_0, X_1 - \mathcal{F} \circ NL(X, u)(T)) \\ Y := e^{T L_1} X_0 + \mathcal{F} \circ L_2 v + \mathcal{F} \circ NL(X, u) \end{cases}$$

□

A simple quadratic obstruction

$$\begin{cases} \dot{x}_1 = u \\ \dot{x}_2 = x_1^2 \end{cases} \quad \dot{x}_2 \geq 0: \text{ no controllability.}$$

A quadratic obstruction in small time

$$\begin{cases} \dot{x}_1 = u \\ \dot{x}_2 = x_1 \\ \dot{x}_3 = x_1^2 - x_2^2 \end{cases} \quad \begin{array}{l} \text{If } x_2(0) = x_2(T) = 0, \int_0^T x_2^2 \leq (T/\pi)^2 \int_0^T \dot{x}_2^2 \\ \text{(Poincaré). If } T \text{ is small, } x_3(T) \geq x_3(0): \text{ no} \\ \text{small-time controllability} \end{array}$$

Another small-time obstruction?

$$\begin{cases} \dot{x}_1 = u \\ \dot{x}_2 = x_1 \\ \dot{x}_3 = x_1^3 + x_2^2 \end{cases} \quad \begin{array}{l} \text{Small-time local controllability... but not if} \\ \text{we ask } |u|_{W^{1,\infty}} \ll 1! \end{array}$$

[Beauchard-Marbach, Quadratic obstructions to small-time local controllability for scalar-input systems, 2018,...]

Previous examples of quadratic obstruction

- The control system

- The case of a non-controllable linearization

Control of the KdV Equation

- KdV Equation

- Quadratic Approximation

- Nonlinear Equation

Control of a Water-Tank

- The Water-Tank System

- (Non)controllability for the Water-Tank

- Kernel for the Quadratic Approximation

- Nonlinear Equation

Conclusion

Previous examples of quadratic obstruction

Schrödinger equation

$$i\partial_t\psi(t,x) = -\partial_x^2\psi(t,x) + u(t)\mu(x)\psi(t,x), \quad x \in (0,1) \text{ with Dirichlet B.C.}$$

Theorem (Smallness of reachable space, Ball, Marsden & Slemrod 1982)

Let $\psi_0 \in L^2(0,1)$. The set

$$\{\psi(T, \cdot) : T > 0, u \in L^2(0,T), \psi \text{ solution with } \psi(0, \cdot) = \psi_0\}$$

is contained in a countable union of compact subsets of $L^2(0,1)$.

Schrödinger equation

$$i\partial_t\psi(t,x) = -\partial_x^2\psi(t,x) + u(t)\mu(x)\psi(t,x), \quad x \in (0,1) \text{ with Dirichlet B.C.}$$

Theorem (Smallness of reachable space, Ball, Marsden & Slemrod 1982)

Let $\psi_0 \in L^2(0,1)$. The set

$$\{\psi(T, \cdot) : T > 0, u \in L^2(0,T), \psi \text{ solution with } \psi(0, \cdot) = \psi_0\}$$

is contained in a countable union of compact subsets of $L^2(0,1)$.

Theorem (Local controllability in H^3 around the ground state, Beauchard & Laurent 2010)

$(\varphi_k)_k$ eigenfunctions of $-\partial_x^2$. If $|\langle \mu\varphi_1, \varphi_k \rangle_{L^2}| \geq ck^{-3}$, for every $T > 0$, for every ψ_0, ψ_1 with appropriate boundary conditions and

$$\|\psi_0 - \varphi_1\|_{H^3} + \|\psi_1 - e^{-i\lambda_1 T}\varphi_1\|_{H^3} \text{ small enough,}$$

there exists $u \in L^2(0,T)$ such that the associated solution satisfies $\psi(T, \cdot) = \psi_1$.

Proof.

Variant of the linear test



Theorem (Quadratic obstruction for small-time local controllability, Coron, Beauchard, Morancey, Bournissou)

If $\langle \mu \varphi_1, \varphi_K \rangle = 0$, under some assumptions on μ , there exists $A > 0$, $T > 0$ and $\eta > 0$ such that for every u with $\|u\|_{H^3(0,T)} < \eta$,

$$\pm \Im \langle \psi(T), \varphi_K e^{-i\lambda_1 T} \rangle \geq A \|u_3\|_{L^2}^2 - C \|\psi(T) - \varphi_1 e^{-i\lambda_1 T}\|_{L^2}^2$$

where $u_0 = u$, $u_{k+1}(t) := \int_0^t u_k(s) ds$.

Theorem (Small-time local controllability with oscillating controls, Bournissou 2022)

Under more assumptions on μ , the Schrödinger equation with bilinear controls is small-time locally controllable around $\varphi_1 e^{-i\lambda_1 T}$ with targets in $D((-\partial_x^2)^{11/2})$ and controls small in $H_0^2(0, T)$.

Proofs.

$$\psi(t, x) = \varphi_1 e^{-i\lambda_1 T} + \psi_{\text{lin}}(u) + \psi_{\text{quad}}(u) + \psi_{\text{cub}}(u) + \text{error}.$$

□

Theorem (Viscous Burgers equation, Marbach 2018)

If $y(0, x) = 0$ and

$$\partial_t y(t, x) - \partial_x^2 y(t, x) + y(t, x) \partial_x y(t, x) = u(t), \quad x \in (0, 1) \text{ with Dirichlet B.C.,}$$

for some test function ρ , $T > 0$ small enough, and $u_1(t) := \int_0^t u(s) ds$,

$$\langle \rho, y(T, \cdot) \rangle \geq k \|u_1\|_{H^{-1/4}}^2.$$

Theorem (Viscous Burgers equation, Marbach 2018)

If $y(0, x) = 0$ and

$$\partial_t y(t, x) - \partial_x^2 y(t, x) + y(t, x) \partial_x y(t, x) = u(t), \quad x \in (0, 1) \text{ with Dirichlet B.C.},$$

for some test function ρ , $T > 0$ small enough, and $u_1(t) := \int_0^t u(s) ds$,

$$\langle \rho, y(T, \cdot) \rangle \geq k \|u_1\|_{H^{-1/4}}^2.$$

Theorem (Nonlinear heat equation, Beauchard et Marbach 2020)

If $\langle \Gamma[0], \varphi_0 \rangle = 0$, under some assumptions on $\Gamma \in C^2(H_N^1; H_N^{-1})$, there exists $A \neq 0$ such that for every $\epsilon > 0$, there exist $T > 0$ and $\eta > 0$ such that for every $\delta \in [-1, 1]$ and $\|u\|_{H^{2n+2}} < \eta$, if

$$\partial_t z(t, x) - \partial_x^2 z(t, x) = u(t) \Gamma(z(t))(x), \quad x \in (0, 1) \text{ with Neuman B.C.},$$

and $z(0) = \delta \varphi_0$ and for $j \geq 1$, $\langle z(T), \varphi_j \rangle \neq 0$,

$$|\langle z(T), \varphi_0 \rangle - \delta + A \|u_n\|_{L^2}^2| \leq \epsilon (|\delta| + \|u_n\|_{L^2}^2).$$

where $u_0 = u$, $u_{k+1}(t) := \int_0^t u_k(s) ds$.

Control of the KdV Equation

KdV equation

$$\begin{cases} \partial_t y + \partial_x y + \partial_x^3 y + y \partial_x y = 0, & (t, x) \in (0, T) \times (0, L) \\ y(t, 0) = y(t, L) = 0, \partial_x y(t, L) = u(t) & t \in (0, T) \end{cases}$$

KdV equation linearized around 0

$$\begin{cases} \partial_t y_1 + \partial_x y_1 + \partial_x^3 y_1 = 0, & (t, x) \in (0, T) \times (0, L) \\ y_1(t, 0) = y_1(t, L) = 0, \partial_x y_1(t, L) = u(t) & t \in (0, T) \end{cases}$$

KdV equation

$$\begin{cases} \partial_t y + \partial_x y + \partial_x^3 y + y \partial_x y = 0, & (t, x) \in (0, T) \times (0, L) \\ y(t, 0) = y(t, L) = 0, \partial_x y(t, L) = u(t) & t \in (0, T) \end{cases}$$

KdV equation linearized around 0

$$\begin{cases} \partial_t y_1 + \partial_x y_1 + \partial_x^3 y_1 = 0, & (t, x) \in (0, T) \times (0, L) \\ y_1(t, 0) = y_1(t, L) = 0, \partial_x y_1(t, L) = u(t) & t \in (0, T) \end{cases}$$

Theorem (Rosier 1997)

The linearized KdV equation is controllable in some time (equivalently in arbitrarily small time) iff $L \notin \mathcal{N} := \left\{ 2\pi \sqrt{\frac{k^2 + kl + l^2}{3}}, (k, l) \in (\mathbb{N}^)^2 \right\}$.*

If $L \in \mathcal{N}$, there is some finite dimensional unreachable space \mathcal{M} .

Theorem (Rosier 1997)

If $L \notin \mathcal{N}$, the nonlinear KdV equation is small-time locally controllable.

Theorem (Coron and Crépeau 2004)

If L can be written in a unique way as $L = 2\pi\sqrt{\frac{k^2+kl+l^2}{3}}$ and that $k = l$, the nonlinear KdV equation is small-time locally controllable.

Theorem (Cerpa 2007, Crépeau and Cerpa 2009)

If $L \in \mathcal{N}$, there exists $T > 0$ such that the nonlinear KdV equation is locally controllable in time T .

Theorem (Rosier 1997)

If $L \notin \mathcal{N}$, the nonlinear KdV equation is small-time locally controllable.

Theorem (Coron and Crépeau 2004)

If L can be written in a unique way as $L = 2\pi\sqrt{\frac{k^2+kl+l^2}{3}}$ and that $k = l$, the nonlinear KdV equation is small-time locally controllable.

Theorem (Cerpa 2007, Crépeau and Cerpa 2009)

If $L \in \mathcal{N}$, there exists $T > 0$ such that the nonlinear KdV equation is locally controllable in time T .

Theorem (Coron K Nguyen 2020)

If $k \neq l \in \mathbb{N}^$, $L = 2\pi\sqrt{\frac{k^2+kl+l^2}{3}}$ and $2k + l \notin 3\mathbb{N}$, lack of small-time local controllability of the nonlinear KdV equation for H^3 initial conditions with controls small in $H^1(0, T)$.*

Order 2

$$\begin{cases} \partial_t y_1 + \partial_x y_1 + \partial_x^3 y_1 = 0, & (t, x) \in (0, T) \times (0, L) \\ y_1(t, 0) = y_1(t, L) = 0, \partial_x y_1(t, L) = u(t) & t \in (0, T) \end{cases}$$

Order 2

$$\begin{cases} \partial_t y_2 + \partial_x y_2 + \partial_x^3 y_2 = -y_1 \partial_x y_1, & (t, x) \in (0, T) \times (0, L) \\ y_2(t, 0) = y_2(t, L) = \partial_x y_2(t, L) = 0 & t \in (0, T) \end{cases}$$

Lemma

If $\dim(\mathcal{M}) = 2$, we identify $\mathcal{M} \approx \mathbb{C}$, and then for some explicit $p \in \mathbb{R}$ and function ϕ .

$$y_{2|\mathcal{M}}(t) = \int_0^L \int_0^t y_1(s, x)^2 e^{ip(t-s)} \phi(x) dx ds.$$

Theorem

If $L = 2\pi\sqrt{\frac{k^2+kl+l^2}{3}}$ with $2k+l \notin 3\mathbb{N}$, if T is small and if u steers y_1 from 0 to 0,

$$y_{2|\mathcal{M}} = \int_0^L \int_0^T y_1(s, x)^2 e^{ip(T-s)} \phi(x) dx ds = EN(u)^2(1 + O(T^{1/4}))$$

where $E \in \mathbb{C} \setminus \{0\}$ and $N(u) \sim \|u\|_{H^{-2/3}}$.

Proof.


- Take Fourier transform in t . For some explicitly computable function $\Lambda(x, z)$,

$$\hat{y}(z, x) = \hat{u}(z)\Lambda(z, x)$$
- Paley-Wiener: if u steers the linearized equation from 0 to 0 then \hat{u} and $\Lambda(\cdot, x)\hat{u}(\cdot)$ are entire and $|\hat{u}(z)| + |\hat{u}(z)\partial_x\Lambda(z, 0)| \leq Ce^{T|\Im(z)|}$.
- Computations $y_{2|\mathcal{M}} = \int \hat{u}(s)\overline{\hat{u}(s-p)}B(s) ds$, $B(s) \underset{s \rightarrow \pm\infty}{\sim} E|s|^{-4/3}$
- In the integral above, the part for $|s| \leq m$ is $\leq CmT^{1/2}\|u\|_{H^{-2/3}}^2$ (we use the Paley-Wiener property here). □

Control of a Water-Tank


The water-tank system

$$\begin{cases} \partial_t H + \partial_x (vH) = 0, & (t, x) \in (0, T) \times (0, L) \\ \partial_t v + \partial_x (gH + v^2/2) = -u(t), & (t, x) \in (0, T) \times (0, L) \\ v(t, 0) = v(t, L) = 0 & t \in (0, T) \\ \ddot{D}(t) = u(t) & t \in (0, T) \end{cases}$$



The water-tank system

$$\begin{cases} \partial_t H + \partial_x(vH) = 0, & (t, x) \in (0, T) \times (0, L) \\ \partial_t v + \partial_x(gH + v^2/2) = -u(t), & (t, x) \in (0, T) \times (0, L) \\ v(t, 0) = v(t, L) = 0 & t \in (0, T) \\ \ddot{D}(t) = u(t) & t \in (0, T) \end{cases}$$



Linearized equation around $H = H_{\text{eq}}, v = 0$

$$\begin{cases} \partial_t h + H_{\text{eq}} \partial_x v = 0, & (t, x) \in (0, T) \times (0, L) \\ \partial_t v + g \partial_x h = -u(t), & (t, x) \in (0, T) \times (0, L) \\ v(t, 0) = v(t, L) = 0 & t \in (0, T) \end{cases}$$

$h(t, L-x) = -h(t, x), v(t, L-x) = v(t, x)$; not controllable. But moving the tank and such that the water is still at the start and end is possible if

$$T > T_* = L / \sqrt{gH_{\text{eq}}}.$$

Theorem (Control using the return method, Coron 2002)

Local controllability in large time: there exists $T > 0, \eta > 0$ such that if

$$\|H_0 - 1\|_{C^1} + \|v_0\|_{C^1} < \eta,$$

$$\|H_1 - 1\|_{C^1} + \|v_1\|_{C^1} < \eta,$$

$$\|D_1 - D_0\| < \eta$$

*then there exists a trajectory such that $H(t=0) = H_0, H(t=T) = H_1,$
 $v(t=0) = v_0, v(t=T) = v_1, D(0) = D_0, D(T) = D_1, \dot{D}(0) = \dot{D}(T) = 0.$*

Theorem (Control using the return method, Coron 2002)

Local controllability in large time: there exists $T > 0, \eta > 0$ such that if

$$\begin{aligned}\|H_0 - 1\|_{C^1} + \|v_0\|_{C^1} &< \eta, \\ \|H_1 - 1\|_{C^1} + \|v_1\|_{C^1} &< \eta, \\ \|D_1 - D_0\| &< \eta\end{aligned}$$

*then there exists a trajectory such that $H(t=0) = H_0, H(t=T) = H_1,$
 $v(t=0) = v_0, v(t=T) = v_1, D(0) = D_0, D(T) = D_1, \dot{D}(0) = \dot{D}(T) = 0.$*

Theorem (Lack of local controllability when the time is not large enough, Coron-K-Nguyen 2021)

For $T < 2T_$, lack of local controllability with controls small in C^0 : there exists $\eta > 0$ such that if $H(t=0) = H(t=T) = H_{\text{eq}}, v(t=0) = v(t=T) = 0,$
 $\dot{D}(0) = \dot{D}(T) = 0,$ and if $\|u\|_{C^0} < \eta,$ then $u = 0.$*

Proof strategy: $(H, v) \approx$ linearized + quadratic, and the quadratic term is $\geq c\|u\|_{H^{-1}}^2.$

Rescaling

$$L = 1, H_{\text{eq}} = 1, g = 1, T_* = 1.$$

Linearized equation

$$\partial_t h_1 + \partial_x v_1 = 0$$

$$\partial_t v_1 + \partial_x h_1 = -u(t)$$

$$v_1(t, 0) = v_1(t, 1) = 0$$

Rescaling

$$L = 1, H_{\text{eq}} = 1, g = 1, T_* = 1.$$

Quadratic term

$$\partial_t h_2 + \partial_x v_2 = -\partial_x (h_1 v_1)$$

$$\partial_t v_2 + \partial_x h_2 = -\partial_x (v_1^2/2)$$

$$v_2(t, 0) = v_2(t, 1) = 0$$

Lemma

$$(h_2(T, \cdot), \phi) + (v_2(T, \cdot), \psi) = \int_{[0, T]^2} K_{T, \phi, \psi}(s_1, s_2) u(s_1) u(s_2) ds_1 ds_2$$

for some explicitly computable kernel $K_{T, \phi, \psi}$.

Formula for the kernel (do not read)

With $\Phi(x) = (\phi(x) + \psi(x))/2$ for $0 < x < 1$ and $(\phi(-x) - \psi(-x))/2$ for $-1 < x < 0$,

$$2K_{T,\phi,\psi}(s_1, s_2) =$$

$$\left\{ \begin{array}{l} \int_{-2T+2s_2}^0 \Phi(s+T-s_2) ds + 2(T-s_2)\Phi(T-s_2) - 4(T-s_2)\Phi(T-s_1) \\ \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \text{if } 2T-1 < s_1+s_2 < 2T \\ \int_{s_2-s_1}^{2-2T+s_2+s_1} \Phi(s-s_2+T) ds + (4T-1-3s_2-s_1)\Phi(T-s_2) - (1+2T-3s_2+s_1)\Phi(T-s_1) \\ \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \text{if } 2T-2 < s_1+s_2 < 2T-1 \\ \int_{2-2T+2s_2}^0 \Phi(s+T-s_2) ds + (1+2T-2s_2)\Phi(T-s_2) - (-1+4T-4s_2)\Phi(T-s_1) \\ \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \text{if } 2T-3 < s_1+s_2 < 2T-2 \\ \int_{s_2-s_1}^{4-2T+s_2+s_1} \Phi(s+T-s_2) ds + (-2+4T-3s_2-s_1)\phi(T-s_2) - (2+2T-3s_2+s_1)\phi(T-s_1) \\ \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \text{if } 2T-4 < s_1+s_2 < 2T-3 \end{array} \right.$$

Lemma

$\Phi(x) = (\phi(x) + \psi(x))/2$ for $0 < x < 1$ and $(\phi(-x) - \psi(-x))/2$ for $-1 < x < 0$. If $1 < T < 2$ and if the control u steers the linearized equation from 0 to 0 (apart from maybe moving the tank),

$$(h_2(T, \cdot), \phi) + (v_2(T, \cdot), \psi) = \int_{[0, T-1]^2} K_{T, \phi, \psi}^{\text{red}}(s_1, s_2) u(s_1) u(s_2) ds_1 ds_2$$

with

$$K_{T, \phi, \psi}^{\text{red}}(s_1, s_2) = \frac{3}{2}(1 - |s_2 - s_1|) (\Phi(T - s_1 \vee s_2) - \Phi(T - s_1 \wedge s_2))$$

Lemma

$\Phi(x) = (\phi(x) + \psi(x))/2$ for $0 < x < 1$ and $(\phi(-x) - \psi(-x))/2$ for $-1 < x < 0$. If $1 < T < 2$ and if the control u steers the linearized equation from 0 to 0 (apart from maybe moving the tank),

$$(h_2(T, \cdot), \phi) + (v_2(T, \cdot), \psi) = \int_{[0, T-1]^2} K_{T, \phi, \psi}^{\text{red}}(s_1, s_2) u(s_1) u(s_2) ds_1 ds_2$$

with

$$K_{T, \phi, \psi}^{\text{red}}(s_1, s_2) = \frac{3}{2}(1 - |s_2 - s_1|) (\Phi(T - s_1 \vee s_2) - \Phi(T - s_1 \wedge s_2))$$

Proposition

Φ 1-periodic, $\Phi(s) = s$ for $s \in [1, T]$. For $1 < T < 2$ and $U(s) = \int_0^s u(s') ds'$

$$(h_2(T, \cdot), \phi) + (v_2(T, \cdot), \psi) \geq 3(2 - T) \|U\|_{L^2(0, T-1)}^2$$

End of the proof.

$$(h, v) \approx \underbrace{(h_1, v_1)}_{\text{linear in } u} + \underbrace{(h_2, v_2)}_{\text{quadratic in } u}$$

□

Conclusion

Quadratic obstruction for small-time local controllability

- Finite-dimensional systems
- Schrödinger equation with bilinear controls
- Viscous Burgers equation
- Some nonlinear heat equations

Quadratic obstruction for small-time local controllability

- Finite-dimensional systems
- Schrödinger equation with bilinear controls
- Viscous Burgers equation
- Some nonlinear heat equations

KdV

- For some critical lengths, lack of small-time local controllability for controls small in H^1 .
- Small-time local controllability with less regular controls?
- Minimal time for local-controllability?

Quadratic obstruction for small-time local controllability

- Finite-dimensional systems
- Schrödinger equation with bilinear controls
- Viscous Burgers equation
- Some nonlinear heat equations

KdV

- For some critical lengths, lack of small-time local controllability for controls small in H^1 .
- Small-time local controllability with less regular controls?
- Minimal time for local-controllability?

Water-tank

- A trajectory which is natural for the water-tank is possible for the linearized equation but not for the nonlinear equation.
- Minimal time for the local-controllability to hold?

That's all folks!

Bonus: Coercivity of an arbitrary scalar product for the water tank

QuestionCoercivity of Q_Ψ :

$$Q_\Psi(u) = \int_{[a,b]^2} u(s_1)u(s_2)(1 + \epsilon|s_2 - s_1|)(\Psi(s_1 \wedge s_2) - \Psi(s_1 \vee s_2)) ds_1 ds_2?$$

(with $\Psi = -\Phi(T - s)$, $Q_\Psi = \langle \Phi, \text{order 2 for the water-tank} \rangle$.)

QuestionCoercivity of Q_Ψ :

$$Q_\Psi(u) = \int_{[a,b]^2} u(s_1)u(s_2)(1 + \epsilon|s_2 - s_1|)(\Psi(s_1 \wedge s_2) - \Psi(s_1 \vee s_2)) ds_1 ds_2?$$

(with $\Psi = -\Phi(T - s)$, $Q_\Psi = \langle \Phi, \text{order 2 for the water-tank} \rangle$.)**Lemma** $\Psi \in C^1$, $\Psi' \geq c > 0$. Then,

$$Q_\Psi(U') \geq \alpha \|U\|_{L^2}^2 \text{ for every } U \in H_0^1(a, b)$$

iff

$$\int_a^b \Psi'(s) ds \int_a^b \frac{1}{\Psi'(s)} ds < (b - a + 2\epsilon^{-1})^2$$

Proof.

Integrate by parts; consider the resulting formula as a quadratic form on $L^2(\Psi'(s) ds)$; see that on a stable space with codimension 2, $Q_\Psi = \text{Identity}$; compute explicitly the 2×2 matrix on the orthogonal and study its positivity. □